


1. A cup contains 250 g of water. Find the total positive charge present in the cup of water.


a) $1.34 \times 10^{19} \text{ C}$
 b) $1.34 \times 10^7 \text{ C}$
 c) $2.43 \times 10^{19} \text{ C}$
 d) $2.43 \times 10^7 \text{ C}$

2. A particle of mass m and charge $-q$ enters the region between the two charged plates initially moving along x -axis with speed v_x as shown in figure. The length of plate is L and a uniform electric field E is maintained between the plates. The vertical deflection of the particle at the far edge of the plate is




a) $\frac{qEL^2}{2mv_x^2}$
 b) $\frac{qEL^2}{2mv_x}$
 c) $\frac{2mv_x^2}{qEL^2}$
 d) $\frac{2mv_x}{qE^2L}$

3. Two point charge $q_1 = -4\mu\text{C}$ and $q_2 = 8\mu\text{C}$ are lying on the y -axis. They are equidistant from the point P , which lies on the x -axis. A small object of charge $q_0 = 8\mu\text{C}$ and mass $m = 12 \text{ g}$ is placed at P . What it is released, what is its acceleration (in m s^{-2}) ? (Neglect the effect of gravity)



a) $3\sqrt{3}\hat{i} + 9\hat{j}$
 b) $9\hat{i} + 3\sqrt{3}\hat{j}$
 c) $3\hat{i} + 3\sqrt{3}\hat{j}$
 d) $3\sqrt{3}\hat{i} + 3\hat{j}$


4. Let σ be the uniform surface charge density of two infinite thin plane sheets shown in figure. Then the electric field in three different region E_I , E_{II} and E_{III} are



Rough Work


- a) $\vec{E}_I = 0, \vec{E}_{II} = \frac{\sigma}{\epsilon_0} \hat{n}, \vec{E}_{III} = 0$
- b) $\vec{E}_I = \frac{2\sigma}{\epsilon_0} \hat{n}, \vec{E}_{II} = 0, \vec{E}_{III} = \frac{2\sigma}{\epsilon_0} \hat{n}$
- c) $\vec{E}_I = -\frac{\sigma}{\epsilon_0} \hat{n}, \vec{E}_{II} = 0, \vec{E}_{III} = \frac{\sigma}{\epsilon_0} \hat{n}$
- d) $\vec{E}_I = \frac{\sigma}{2\epsilon_0} \hat{n}, \vec{E}_{II} = 0, \vec{E}_{III} = \frac{\sigma}{2\epsilon_0} \hat{n}$

5. A finite ladder is constructed by connecting several sections of 2μ , $4\mu F$ capacitor combinations as shown in figure. It is terminated by a capacitor of capacitance C. The value that should be chosen for C, such that the equivalent capacitance of the ladder between the points A and B becomes independent of the number of sections in between, is μF .




- a) $4\mu F$
- b) $5\mu F$
- c) $6\mu F$
- d) $7\mu F$

6. A capacitor is made of a flat plate of area A and a second plate having a stair-like structure as shown in figure. If the area of each stair is $\frac{A}{3}$ and the height is d, the capacitance of the arrangement is



- a) $\frac{11\epsilon_0 A}{20d}$
- b) $\frac{18\epsilon_0 A}{11d}$
- c) $\frac{11\epsilon_0 A}{18d}$
- d) $\frac{13\epsilon_0 A}{17d}$


7. In the shown figure, bridge is balanced, the current flowing through 2Ω resistance is



Rough Work


- a) $\frac{10}{7} A$
- b) $\frac{11}{7} A$
- c) $\frac{13}{7} A$
- d) $\frac{8}{7} A$

8. A 100 W bulb B_1 and two 60 W bulbs B_2 and B_3 , are connected to a 250 V source, as shown in figure. Now W_1 , W_2 and W_3 are the output powers of the bulbs B_1 , B_2 and B_3 , respectively. Then




- a) $W_1 > W_2 = W_3$
- b) $W_1 > W_2 > W_3$
- c) $W_1 < W_2 = W_3$
- d) $W_1 < W_2 < W_3$

9. The terminal voltage of the battery, whose emf is 10 V and internal resistance 1Ω , when connected through an external resistance of 4Ω as shown in the figure is



- a) 4 V
- b) 6 V
- c) 8 V
- d) 10 V

10. The value of unknown resistance (x) for which the potential difference between B and D will be zero in the arrangement shown, is




- a) 42Ω
- b) 9Ω
- c) 6Ω
- d) 3Ω

11. An element of $0.05\hat{i}$ m is placed at the origin as shown in figure which carries a


Rough Work

large current of 10 A. The magnetic field at a distance of 1 m in perpendicular direction is




- a) $4.5 \times 10^{-8} \text{ T}$
- b) $5.5 \times 10^{-8} \text{ T}$
- c) $5.0 \times 10^{-8} \text{ T}$
- d) $7.5 \times 10^{-8} \text{ T}$

12. In the given circuit, a galvanometer with a resistance of 70Ω is converted to an ammeter by a shunt resistance of 0.05Ω , total current measured by this device is




- a) 0.88 A
- b) 0.77 A
- c) 0.55 A
- d) 0.99 A

13. A particle of charge q and mass m is projected with a velocity v_0 towards a circular region having uniform magnetic field B perpendicular and into the plane of paper, from point P as shown in figure. R is the radius and O is the centre of the circular region. If the line OP makes an angle θ with the direction of v_0 then the value of v_0 so that particle passes through O is




- a) $\frac{qBR}{m \sin \theta}$
- b) $\frac{qBR}{2m \sin \theta}$
- c) $\frac{2qBR}{m \sin \theta}$
- d) $\frac{3qBR}{2m \sin \theta}$

14. A thick current carrying cable of radius ' R ' carries current ' I ' uniformly distributed across its cross-section. The variation of magnetic field $B(r)$ due to the cable with the distance ' r ' from the axis of the cable is represented by



15. A helmholtz coil has a pair of loops, each with N turns and radius R . They are placed coaxially at distance R and the same current I flows through the loops in the same direction. The magnitude of magnetic field at P , midway between the centre A and C , is given by



a) $\frac{4N\mu_0 I}{5^{1/2} R}$

b) $\frac{4N\mu_0 I}{5^{3/2} R}$

c) $\frac{8N\mu_0 I}{5^{3/2} R}$

d) $\frac{8N\mu_0 I}{5^{1/2} R}$

16. A short bar magnet placed with its axis at 30° with a uniform external magnetic field of 0.35 T experiences a torque of magnitude equal to $4.5 \times 10^{-2} \text{ N m}$. The magnitude of magnetic moment of the given magnet is


a) 26 J T^{-1}

b) 2.6 J T^{-1}

c) 0.26 J T^{-1}

d) 0.026 J T^{-1}

17. Some equipotential surfaces of the magnetic scalar potential are shown in figure. Magnetic field at a point in the region (in T) is



a) 10^{-4}


b) 0.5×10^{-4}

c) 2×10^{-4}

d) None


Rough Work

18. In a uniform magnetic field of 0.049 T, a magnetic needle performs 20 complete oscillations in 5 seconds as shown. The moment of inertia of a needle is $9.8 \times 10^{-6} \text{ kg m}^2$. If the magnitude of magnetic moment of the needle is $x \times 10^{-5} \text{ Am}^2$; then the value of 'x' is




- a) $5\pi^2$
 b) $128\pi^2$
 c) $50\pi^2$
 d) $1280\pi^2$
19. A susceptibility of a certain magnetic material is 400. What is the class of the magnetic material ?
- a) Ferromagnetic
 b) Diamagnetic
 c) Ferroelectric
 d) Paramagnetic
20. Two identical bar magnets are fixed with their centres at a distance d apart. a stationary charge Q is placed at P in between the gap

of the two magnets at a distance D from the centre O as shown in the figure. The force on the charge Q is




- a) zero
 b) directed along OP
 c) directed along PO
 d) directed perpendicular to the plane of paper
21. A voltmeter which can measure 2 V is constructed by using a galvanometer of resistance 12Ω and that produces maximum deflection for the current of 2mA, then the resistance R is




- a) 888Ω
 b) 988Ω
 c) 898Ω
 d) 999Ω

22. A triangular shaped wire carrying 10 A current is placed in a uniform magnetic field of 0.5 T, as shown in figure. The magnetic force on segment CD is
(Given $BC = CD = BD = 5$ cm.)



- a) 0.126 N
b) 0.312 N
c) 0.216 N
d) 0.245 N
23. A 7 V battery with internal resistance 2Ω and a 3 V battery with internal resistance 1Ω are connected to a 10Ω resistor as shown in figure, the current in 10Ω resistor is




- a) 0.27 A
b) 0.31 A
c) 0.031 A
d) 0.53 A
24. A silver wire has temperature coefficient of resistivity $4 \times 10^{-3} \text{ } ^\circ\text{C}^{-1}$ and its resistance at 20

$^\circ\text{C}$ is 10Ω . Neglect any change in dimensions due to the change in temperature, its resistance at $40 \text{ } ^\circ\text{C}$ is

- a) 0.8Ω
b) 1.8Ω
c) 9.2Ω
d) 10.8Ω


25. Suppose the drift velocity v_d in a material varied with the applied electric field E as $v_d \propto \sqrt{E}$. Then V-I graph for a wire made of such a material is best given by



26. In an atom electrons revolves around the nucleus along a path of radius $0.72 \text{ \AA}$ making 9.4×10^{18} revolution per second. The equivalent current is ($e = 1.6 \times 10^{-19} \text{ C}$)


a) 1.2 A
b) 1.5 A
c) 1.4 A
d) 1.8 A

27. When a potential difference of 10^3 V is applied between A and B, a charge of 0.75 mC is stored in the system of capacitors as shown. The value of C is (in μF) and energy stored (in J) in the equivalent capacitor is



a) $\frac{1}{2}, 2$
b) 2, 1
c) 2.5, 2
d) 3, 1

28. The charge on $3 \mu\text{F}$ capacitor shown in the figure is




a) $2 \mu\text{C}$
b) $10 \mu\text{C}$
c) $6 \mu\text{C}$
d) $8 \mu\text{C}$

29. The electric field components in the given


figure are $E_x = \alpha x^{1/2}$, $E_y = E_z = 0$ in which

$\alpha = 800 \text{ NC}^{-1} \text{ m}^{-1/2}$. The charge within the cube if net flux in (in 10^{-12} C) through the cube is $1.05 \text{ N m}^2 \text{ C}^{-1}$, is approx (assume $a = 0.1 \text{ m}$)






a) 9.27
b) 8.27
c) 5.97
d) 4.97


30. Which of the following curves represent electric field lines correctly ?



Rough Work



31. Consider the two figures given below.



Which of the following statements regarding the experiment is true?

- a) The solubility of a gas in liquid in beaker (i) is greater than that in beaker (ii).
- b) The solubility of a gas in beaker (i) is less than that in beaker (ii).
- c) The solubility of a gas is equal in both beakers.
- d) The solubility of a gas remains unaffected by change in weights.

32. The value of Henry's law constant for some gases at 293 K is given below. Arrange the gases in the increasing order of their solubility.

He: 144.97 kbar, H_2 : 69.16 kbar,

N_2 : 76.48 kbar, O_2 : 34.86 kbar


(a) $He < N_2 < H_2 < O_2$

(b) $O_2 < H_2 < N_2 < He$

(c) $H_2 < N_2 < O_2 < He$

(d) $He < O_2 < N_2 < H_2$

33.



X, Y and Z in the above graph are


a) $X = p_1 + p_2$, $Y = 1$, $Z = 0$

b) $X = p_1 + p_2$, $Y = 0$, $Z = 1$

c) $X = p_1 \times p_2$, $Y = 0$, $Z = 1$


d) $X = p_1 - p_2$, $Y = 1$, $Z = 0$

34. Study the figures given below and mark the correct statement.



- a) (i) Nitric acid + Water,
 (ii) Acetone + Ethyl alcohol
 b) (i) Water + Ethyl alcohol,
 (ii) Acetone + Benzene
 c) (i) Acetone + Ethyl alcohol,
 (ii) Acetone + Chloroform
 d) (i) Benzene + Chloroform,
 (ii) Acetone + Chloroform

35. If semipermeable membrane is placed between the solvent and solution as shown in the given figure then



- a) the solvent molecules will flow through the membrane from solution to pure solvent
 b) the solvent molecules will flow continuously till the equilibrium is attained
 c) the flow of the solvent from its side to solution side across a semipermeable membrane can be stopped if some extra pressure (called osmotic pressure) is applied on the solution.
 d) both (b) and (c).

36. Why is the molecular mass determined by measuring colligative property in case of some solutes is abnormal?

- a) Due to association or dissociation of solute molecules.
 b) Due to insolubility of solute molecules.
 c) Due to decomposition of solute molecules.
 d) Due to large size of solute molecules.

37. If α is the degree of dissociation of Na_2SO_4 , the van't Hoff's factor (i) used for calculating the molecular mass is

- a) $1 + \alpha$
 b) $1 - \alpha$
 c) $1 + 2\alpha$
 d) $1 - 2\alpha$

38. Elevation in the boiling point for 1 molal solution of glucose is 2 K. The depression in the freezing point for 2 molal solution of glucose in the same solvent is 2K. The relation between K_b and K_f is
- $K_b = 1.5 K_f$
 - $K_b = 0.5 K_f$
 - $K_b = 2 K_f$
 - $K_b = K_f$
39. Which of the following is the correct cell representation for the given cell reaction?
- $$\text{Zn} + \text{H}_2\text{SO}_4 \rightarrow \text{ZnSO}_4 + \text{H}_2$$
- $\text{Zn} | \text{Zn}^{2+} || \text{H}^+ | \text{H}_2$
 - $\text{Zn} | \text{Zn}^{2+} || \text{H}^+, \text{H}_2 | \text{Pt}$
 - $\text{Zn} | \text{ZnSO}_4 || \text{H}_2\text{SO}_4 | \text{Zn}$
 - $\text{Zn} | \text{H}_2\text{SO}_4 || \text{ZnSO}_4 | \text{H}_2$
40. The cell reaction of the galvanic cell :
- $$\text{Cu}(s) | \text{Cu}_{(\text{aq})}^{2+} || \text{Hg}_{(\text{aq})}^{2+} | \text{Hg}(l)$$
- $\text{Hg} + \text{Cu}^{2+} \rightarrow \text{Hg}^{2+} + \text{Cu}$
 - $\text{Hg} + \text{Cu}^{2+} \rightarrow \text{Cu}^+ + \text{Hg}^+$
 - $\text{Cu} + \text{Hg} \rightarrow \text{CuHg}$
 - $\text{Cu} + \text{Hg}^{2+} \rightarrow \text{Cu}^{2+} + \text{Hg}$
41. The number of electrons delivered at the cathode during electrolysis by a current of 1 ampere in 60 seconds is
- (charge on electron = 1.6×10^{-19})
- 6×10^{23}
 - 6×10^{20}
 - 3.75×10^{20}
 - 7.48×10^{23}
42. The molar conductance of NaCl, HCl and CH_3COONa at infinite dilution are 126.45, 426.16 and 91.0 $\text{S cm}^2 \text{mol}^{-1}$ respectively. The molar conductance of CH_3COOH at Infinite dilution is. Choose the right option for your answer.
- 540.48 $\text{S cm}^2 \text{Mol}^{-1}$
 - 201.28 $\text{S cm}^2 \text{Mol}^{-1}$
 - 390.71 $\text{S cm}^2 \text{Mol}^{-1}$
 - 698.28 $\text{S cm}^2 \text{Mol}^{-1}$
43. The standard electrode potential (E°) for OCl^-/Cl^- and $\text{Cl}^- / \frac{1}{2} \text{Cl}_2$ respectively are 0.94 V and -1.36 V. The E° value for $\text{OCl}^- / \frac{1}{2} \text{Cl}_2$ will be
- 0.42 V
 - 2.20 V
 - 0.52 V
 - 1.04 V

44. Which cell will measure standard electrode potential of copper electrode ?

- a) $\text{Pt}_{(s)} | \text{H}_2(\text{g}, 0.1 \text{ bar}) | \text{H}^+(\text{aq}, 1 \text{ M}) || \text{Cu}^{2+}(\text{aq}, 1 \text{ M}) | \text{Cu}$
 b) $\text{Pt}_{(s)} | \text{H}_2(\text{g}, 1 \text{ bar}) | \text{H}^+(\text{aq}, 1 \text{ M}) || \text{Cu}^{2+}(\text{aq}, 2 \text{ M}) | \text{Cu}$
 c) $\text{Pt}_{(s)} | \text{H}_2(\text{g}, 1 \text{ bar}) | \text{H}^+(\text{aq}, 1 \text{ M}) || \text{Cu}^{2+}(\text{aq}, 1 \text{ M}) | \text{Cu}$
 d) $\text{Pt}_{(s)} | \text{H}_2(\text{g}, 1 \text{ bar}) | \text{H}^+(\text{aq}, 0.1 \text{ M}) || \text{Cu}^{2+}(\text{aq}, 1 \text{ M}) | \text{Cu}$

45. For a cell reaction : $M_{(aq)}^{n+} + ne^- \rightarrow M_{(s)}$, the Nernst equation for electrode potential at any concentration measured with respect to standard hydrogen electrode is represented as

- a) $E_{(M^{n+}/M)} = E_{(M^{n+}/M)}^{\circ} - \frac{RT}{nF} \ln \frac{1}{[M^{n+}]}$
 b) $E_{(M/M^{n+})} = E_{(M/M^{n+})}^{\circ} - \frac{RT}{nF} \ln \frac{[M^{n+}]}{[M]}$
 c) $E_{(M^{n+}/M)} = E_{(M^{n+}/M)}^{\circ} - \frac{RT}{nF} \log \frac{1}{[M]}$
 d) $E_{(M^{n+}/M)} = E_{(M^{n+}/M)}^{\circ} - \frac{RT}{nF} \ln [m^{n+}]$

46. Molar conductivity 0.025 mol L^{-1} methanoic acid is $46.1 \text{ S cm}^2 \text{ mol}^{-1}$, the degree of dissociation and dissociation constant will be

(Give $\lambda_{\text{H}^+}^{\circ} = 349.6 \text{ S cm}^2 \text{ mol}^{-1}$ and

$\lambda_{\text{HCOO}^-}^{\circ} = 54.6 \text{ S cm}^2 \text{ mol}^{-1}$)


- a) 11.4%, $3.67 \times 10^{-4} \text{ mol L}^{-1}$
 b) 22.8%, $1.83 \times 10^{-4} \text{ mol L}^{-1}$
 c) 52.2%, $4.25 \times 10^{-4} \text{ mol L}^{-1}$
 d) 1.14%, $3.67 \times 10^{-6} \text{ mol L}^{-1}$

47. Match the rate law given in column I with the dimensions of rate constants given in column II and mark the appropriate choice

Column I	Column II
(A) Rate = $k[\text{NH}_3]^0$	(i) $\text{mol L}^{-1} \text{s}^{-1}$
(B) Rate = $k[\text{H}_2\text{O}_2][\text{I}^-]$	(ii) $\text{L mol}^{-1} \text{s}^{-1}$
(C) Rate = $k[\text{CH}_3\text{CHO}]^{3/2}$	(iii) s^{-1}
(D) Rate = $k[\text{C}_2\text{H}_5\text{Cl}]$	(iv) $\text{L}^{1/2} \text{mol}^{-1/2} \text{s}^{-1}$

- a) (A) $\rightarrow$ (iv), (B) $\rightarrow$ (iii), (C) $\rightarrow$ (ii), (D) $\rightarrow$ (i)
 b) (A) $\rightarrow$ (i), (B) $\rightarrow$ (ii), (C) $\rightarrow$ (iii), (D) $\rightarrow$ (iv)
 c) (A) $\rightarrow$ (ii), (B) $\rightarrow$ (i), (C) $\rightarrow$ (iv), (D) $\rightarrow$ (iii)
 d) (A) $\rightarrow$ (i), (B) $\rightarrow$ (ii), (C) $\rightarrow$ (iv), (D) $\rightarrow$ (iii)

48. Observe the given graphs carefully.




Which of the given orders are shown by the graphs respectively ?

- | (i) | (ii) |
|-----------------|-------------|
| a) Zero order | First order |
| b) First order | Zero order |
| c) First order | First order |
| d) Second order | Zero order |


49. Match the plots in column I with their orders in column II and mark the appropriate choice.

column I


Column II




(i) Zero order



(ii) First order







(ii) Second



(iv) Third order

- a) (A) → (iii), (B) → (ii), (C) → (i), (D) → (iv)
 b) (A) → (i), (B) → (ii), (C) → (iii), (D) → (iv)
 c) (A) → (iv), (B) → (iii), (C) → (ii), (D) → (i)
 d) (A) → (ii), (B) → (i), (C) → (iii), (D) → (iv)

50. According to Arrhenius equation, rate constant k is equal to $Ae^{-E_a/RT}$. Which of the following options represents the graph of $\ln k$ vs $\frac{1}{T}$?



Rough Work





51. Which of the following statements is not correct for the catalyst ?

- a) It catalyses the forward and backward reaction to the same extent.
- b) It alters ΔG of the reaction.
- c) It is a substance that does not change the equilibrium constant of a reaction
- d) It provides an alternate mechanism by reducing activation energy between reactants and products.

52. Which option is valid for a zero order reaction ?

- a) $t_{3/4} = \frac{3}{2}t_{1/2}$
- b) $t_{1/2} = \frac{4}{2}t_{3/4}$
- c) $t_{1/2} = 2t_{3/4}$
- d) $t_{1/2} = 2t_{3/4}$

53. The variation of the rate of an enzyme catalyzed reaction with substrate concentration is correctly represented by graph



54. Match the column I with column II and mark the appropriate choice.

Column I	Column II
(A) $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	(i) Green
(B) $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$	(ii) Light pink
(C) $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$	(iii) Pale green
(D) $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$	(iv) Pink
(E) Cu_2Cl_2	(v) Colourless

Rough Work

- (a) $(A) \rightarrow (iii), (B) \rightarrow (iv), (C) \rightarrow (i), (D) \rightarrow (ii), (D) \rightarrow (v)$
 (b) $(A) \rightarrow (ii), (B) \rightarrow (iii), (C) \rightarrow (iv), (D) \rightarrow (i), (D) \rightarrow (v)$
 (c) $(A) \rightarrow (v), (B) \rightarrow (ii), (C) \rightarrow (iii), (D) \rightarrow (iv), (D) \rightarrow (i)$
 (d) $(A) \rightarrow (iii), (B) \rightarrow (i), (C) \rightarrow (ii), (D) \rightarrow (iv), (D) \rightarrow (v)$

55. Which of the following compounds is used as the starting material for the preparation of potassium dichromate ?

- (a) $K_2SO_4 \cdot Cr_2(SO_4)_3 \cdot 24H_2O$ (Chrome alum)
 (b) $PbCrO_4$ (Chromite yellow)
 (c) $FeCr_2O_4$ (Chromite)
 (d) $PbCrO_4 \cdot PbO$ (Chrome red)

56. Which of the following reactions is not correct ?

- (a) $MnO_4^- + 8H^+ + 5Fe^{2+} \rightarrow 5Fe^{3+} + Mn^{2+} + 4H_2O$
 (b) $2MnO_2 + 4KOH + O_2 \rightarrow 4KMnO_4 + 2H_2O$
 (c) $2Na_2CrO_4 + 2H^+ \rightarrow Na_2Cr_2O_7 + 2Na^+ + H_2O$
 (d) $K_2Cr_2O_7 + 7H_2SO_4 + 6KI \rightarrow$
 $4K_2SO_4 + Cr_2(SO_4)_3 + 3I_2 + 7H_2O$

57. $A \xrightarrow[green]{4KOH, O_2} 2B + 2H_2O$

$3B \xrightarrow[purple]{4HCl} 2C + MnO_2 + 2H_2O$

$2C \xrightarrow{H_2O, KI} 2A + 2KOH + D$

In the above sequence of reactions, A and D, respectively are

- a) KI and $KMnO_4$
 b) KIO_3 and MnO_2
 c) KI and K_2MnO_4
 d) MnO_2 and KIO_3

58. Given below are two statements.

Statement I : Iron (III) catalyst, acidified $K_2Cr_2O_7$ and neutral $KMnO_4$ have the ability to oxidise I^- to I_2 independently.

Statement II : Manganate ion is

paramagnetic in nature and involves $p\pi - p\pi$ bonding.

In the light of the above statements, choose the correct answer from the options given below

- a) Both statement I and statement II are true
 b) Both statement I and statement II are false
 c) Statement I is true but statement II is false
 d) Statement I is false but statement II is true.

59. Highest oxidation state of manganese in fluorides is +4 (MnF_4) but highest oxidation state in oxides is +7 (Mn_2O_7) because

- a) fluorine is more electronegative than oxygen
 b) fluorine does not possess d-orbitals
 c) fluorine stabilises lower oxidation state

d) in covalent compounds, fluorine can form single bond only while oxygen forms double bond.

60. Strong reducing and oxidising agents among the following, respectively, are

- a) Ce^{4+} and Eu^{2+}
- b) Eu^{2+} and Ce^{4+}
- c) Ce^{3+} and Ce^{4+}
- d) Ce^{4+} and Tb^{4+}

61. The stamens represent:

- (a) microsporangia
- (b) male gametophyte
- (c) male gametes
- (d) microsporophylls

62. Sporopollenin is a constituent of pollen exine. It can be degraded by the action of:

- (a) enzymes
- (b) high temperature
- (c) strong acids
- (d) none of these

63. Which of the following causes chronic respiratory disorders due to pollen allergy?

- (a) *Crotalaria*
- (b) *Eichhornia*
- (c) *Zostera*
- (d) *Parthenium*

64. Feathery stigma occurs in:

- (a) pea
- (b) maize
- (c) *Datura*
- (d) *Hydrilla*

65. Spermiation is the process of the release of sperms from

- (a) seminiferous tubules
- (b) vas deferens
- (c) epididymis
- (d) prostate gland

66. Factors that help in spermiogenesis are secreted by:

- (a) FSH
- (b) GH
- (c) STH
- (d) ATH

67. 1st polar body is formed after which stage of oogenesis?

- (a) 1st meiosis
- (b) 2nd mitosis
- (c) 1st mitosis
- (d) Differentiation

68. Which part of ovary in mammal acts as an endocrine gland after ovulation

- (a) Stroma
- (b) Germinal epithelium
- (c) Vitelline membrane
- (d) Graafian follicle

69. Extrusion of second polar body from egg occurs:

- (a) simultaneously with first cleavage
- (b) after entry of sperm but before fertilisation
- (c) after fertilization
- (d) before entry of sperm

70. Multiload 375 is intrauterine device which contains.

- (a) manganese

- (b) iron
- (c) copper
- (d) calcium

71. Emergency contraceptives could be used to avoid possible pregnancy due to

- (a) Casual unprotected intercourse
- (b) rape
- (c) MTP
- (d) both (a) and (b)

72. Which of the following is a full proof method of contraception?

- (a) Implants
- (b) Lactational amenorrhea
- (c) Condoms
- (d) Sterilisation

73. Hepatitis B is transmitted through

- (a) blood transfusion
- (b) infected mother to fetus
- (c) sexual intercourse
- (d) All of these

- 74. In case of a couple where the male is having a very low sperm count, which technique will be suitable for fertilization?**
- (a) Gamete intracytoplasmic fallopian transfer
 - (b) Artificial Insemination
 - (c) Intracytoplasmic sperm injection
 - (d) Intra uterine transfer
- 75. From the sexually transmitted disease mentioned below, identify the one which does not specifically affect the sex organs?**
- (a) Syphilis
 - (b) Aids
 - (c) Gonorrhea
 - (d) Genital warts
- 76. ABO blood grouping in human beings cites the example of:**
- (a) incomplete dominance
 - (b) co-dominance
 - (c) multiple allelism
 - (d) both (b) and (c)
- 77. Experimental verification of 'chromosomal theory' of inheritance was done by**
- (a) Sutton and Boveri
 - (b) Morgan et al
 - (c) Henking
 - (d) Karl Correns
- 78. Genes with multiple phenotypic effects are known as:**
- (a) epistatic genes
 - (b) duplicate genes
 - (c) pleiotropic genes
 - (d) polygenes
- 79. Which of the following is not an example of recessive autosomal disease?**
- (a) Haemophilia
 - (b) Thalassemia
 - (c) Phenylketonuria
 - (d) Sickle cell anaemia
- 80. Father of a child is colourblind and mother is carrier for colourblindness, the probability of the child being colourblind is:**
- (a) 25%
 - (b) 50%

(c) 100%

(d) 75%

81. Which amino acid is substituted in sickle cell anaemia?

(a) Glutamic acid by valine in α -chain

(b) Glutamic acid by valine in β -chain

(c) Valine by glutamic acid in β -chain

(d) Valine by glutamic acid in α -chain

82. ZZ/ZW type of sex determination is seen in:

(a) platypus

(b) snails

(c) cockroach

(d) peacock

83. If the sequence of bases in coding strand of DNA is ATTCGATG, then the sequence of bases in mRNA will be

(a) TAAGCTAC

(b) UAAGCUAC

(c) ATTCGATG

(d) AUUCGAUG

84. Polycistronic messenger RNA (mRNA)

usually occurs in:

(a) bacteria

(b) prokaryotes

(c) eukaryotes

(d) both (a) and (b)

85. The sequence of structural genes in lac operon is:

(a) lac a, lac y, lac z

(b) lac a, lac z, lac y

(c) lac y, lac z, lac a

(d) lac z, lac y, lac a

86. Which of the following cannot act as inducer?

(a) Glucose

(b) Lactose

(c) Galactose

(d) Both (a) and (c)

87. Which was the last human chromosome to be completely sequenced?

(a) Chromosome 1

(b) Chromosome 11

(c) Chromosome 21

(d) Chromosome X

88. Identify the possible anticodon of Arg from the given sequence.

(a) AGC

(b) GCA

(c) UCA

(d) UGC

89. Which part of the sperm plays an important role in penetrating the egg membrane:

(a) Neck

(b) Tail

(c) Middle piece

(d) Acrosome

90. hCG, hPL and relaxin are produced in women

(a) at the time of puberty

(b) only during pregnancy

(c) at the time of menopause

(d) during menstruation

91. In a certain code, "PLANET" is written as "OMZMDZ" and "ORBIT" is written as "NQSAZ". How is "SYSTEM" written in that code?

a) RXRSDX

b) RYRTDN

c) RXRRDZ

d) RZRTDZ

92. Pointing to a photograph, a man said, "I have no brothers and sisters, but the man's father in the photo is my father's son." How is the man in the photograph related to the speaker?

a) Son

b) Nephew

c) Cousin

d) Self

93. A man starts from point P and walks 5 km east, then takes a left turn and walks 8 km, then takes another left turn and walks 12 km. Finally, he takes a left turn and walks 8 km. How far is he from point P?

a) 2 km

b) 3 km

c) 4 km

d) 5 km

94. In a class of 45 students, A ranks 12th from the top, and B ranks 8th from the bottom. How many students are there between A and B?

a) 24

b) 25

c) 26

d) 27

95. A large cube is painted on two opposite faces and then cut into 216 smaller cubes of equal size. How many cubes have no faces painted?

a) 64

b) 100

c) 96

d) 120

96. If 1st January 2021 was a Friday, what day of the week was 1st January 2025?

a) Tuesday

b) Wednesday

c) Thursday

d) Friday

97. A rectangular grid is made by dividing a square into 5 rows and 5 columns of equal size, forming smaller squares. How many total squares (of all possible sizes) are there in this grid?

a) 25

b) 28

c) 30

d) 55

98. Some chairs are tables.

1. All tables are furniture.

Conclusions:

I. Some chairs are furniture.

II. All chairs are furniture.

a) Only conclusion I follows

b) Only conclusion II follows

c) Both conclusions I and II follow

d) Neither conclusion I nor II follows

99. Mountain : Valley :: Convex : ?

a) Circle

b) Concave

c) Plane

d) Line

100. Choose the odd one out:

- a) Pyramid
- b) Cylinder
- c) Sphere
- d) Cone

Rough Work

ROUGH WORK

[illegible]

ROUGH WORK

[illegible]